ED OBSERVATION UNIT: ANAPHYLAXIS AND ANGIOEDEMA PROTOCOL NYC H+H KINGS COUNTY HOSPITAL CENTER

General Observation Guidelines apply for all ED observation patients.

INCLUSION CRITERIA	EXCLUSION CRITERIA
 Local skin eruptions Clinical impression of angioedema or anaphylaxis Able to speak in full sentences Administration of subcutaneous epinephrine 	 Clinical suspicion or fiberoptic scope evidence of deep airway involvement Requirement for continuous IV epinephrine Respiratory distress SaO2 < 90% (on room air) Tachypnea Stridor or other evidence of impending airway compromise

INTERVENTIONS	OPTIONAL INTERVENTIONS
 Cardiac monitor and oxygen	 Supplemental oxygen Albuterol +/- ipratropium Chest X-ray Imaging Epinephrine auto-injector teaching
saturation monitoring IV Fluids as needed Antihistamines Corticosteroids (IV, PO) Patient education	and prescription

DISPOSITION	
 Home: Resolution or improvement in local skin irritations and/or respiratory function 	 Admission: Significant respiratory symptoms persist Delayed reaction or reoccurrence Does not meet discharge criteria after observation period

Last updated 7/19/2022 Reviewed by Bobak Zonnoor MD, Raymond Beyda MD, Rajesh Verma, MD

ED OBSERVATION UNIT: ANAPHYLAXIS AND ANGIOEDEMA PROTOCOL NYC H+H KINGS COUNTY HOSPITAL CENTER

<u>Sources</u>

- Sampson HA, Muñoz-Furlong A, Campbell RL, et al. Second symposium on the definition and management of anaphylaxis: summary report--second National Institute of Allergy and Infectious Disease/Food Allergy and Anaphylaxis Network symposium. Ann Emerg Med 2006;47(4):373–80.
- 2. Moellman JJ, Bernstein JA, Lindsell C, et al. A consensus parameter for the evaluation and management of angioedema in the emergency department. Acad Emerg Med 2014;21(4):469–84.
- 3. Singer E, Zodda D. Allergy And Anaphylaxis: Principles Of Acute Emergency Management. Emerg Med Pract 2015;17(8):1–19; quiz 20.