

# ED OBSERVATION UNIT: ASTHMA GUIDELINES NYC H+H KINGS COUNTY HOSPITAL CENTER

*General Observation Guidelines apply for all ED observation patients.*

| INCLUSION CRITERIA                                                                                                                                                                                                                              |                                                                                                                                                                                                                                                                                                                                         |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none"> <li>• History of asthma</li> <li>• Initial treatment (nebulizers, steroids, magnesium) and intermediate response (improvement but still wheezing)</li> <li>• Able to speak full phrases/sentences</li> </ul> |                                                                                                                                                                                                                                                                                                                                         |
| EXCLUSION CRITERIA*                                                                                                                                                                                                                             |                                                                                                                                                                                                                                                                                                                                         |
| Hemodynamic instability                                                                                                                                                                                                                         | <ul style="list-style-type: none"> <li>• <math>O_2 &lt; 92\%</math>, HR <math>&gt; 120</math>, RR <math>&gt; 30</math>, SBP <math>&lt; 90</math> mmHg</li> <li>• Pulsus paradoxus <math>&gt; 25</math> mmHg</li> </ul>                                                                                                                  |
| Exam                                                                                                                                                                                                                                            | <ul style="list-style-type: none"> <li>• Absent breath sounds (silent chest)</li> <li>• Change in mental status - agitation, anxiety, lethargy, drowsy, confused</li> <li>• Unable to speak sentences or phrases</li> <li>• Accessory muscle use</li> <li>• Inability to lie in supine position</li> <li>• Cyanosis</li> </ul>          |
| Testing                                                                                                                                                                                                                                         | <ul style="list-style-type: none"> <li>• Peak expiratory flow rate <math>&lt; 40\%</math> of baseline or predicted**</li> <li>• Hypercapnia - <math>PaCO_2 &gt; 45</math> mmHg on VBG</li> <li>• Radiographic evidence of complication requiring inpatient treatment (ie, PTX, PNA)</li> <li>• Cardiac dysrhythmia (ie, SVT)</li> </ul> |
| ER Interventions                                                                                                                                                                                                                                | <ul style="list-style-type: none"> <li>• Mechanical or NIPPV***</li> <li>• Epinephrine or terbutaline (excluding pre-hospital)</li> </ul>                                                                                                                                                                                               |
| Other                                                                                                                                                                                                                                           | <ul style="list-style-type: none"> <li>• Any other need for inpatient admission</li> <li>• Any factor that will preclude discharge in 48 hours</li> </ul>                                                                                                                                                                               |

\* Criteria extrapolated from Milliman admission guidelines and the National Heart, Lung, and Blood Institute's description of severe asthma and high risk features of imminent respiratory failure.<sup>1-3</sup>

\*\*Refer to Mdcalc.com or Table 1 if height not available

\*\*\*The use of NIPPV in asthma is not standard care and is lacking in high quality evidence.<sup>4-5</sup> There is practice variation among ER providers and therefore whether or not a patient was placed on NIPPV should **not independently** rule out or rule in a severe asthma exacerbation. Please refer to exclusion criteria.

Last updated 7/1/2019

Authored by E.Tang MD

Revised by T. Ahmad MD, R. Balakrishnan MD, A. Cai, MD, and S. Brewster MD

**ED OBSERVATION UNIT:**  
**ASTHMA GUIDELINES**  
**NYC H+H KINGS COUNTY HOSPITAL CENTER**

***INTERVENTIONS***

- Bronchodilator nebulizers treatments q2-q4h
- Steroids
- Supplemental O2 prn
- Serial peak flow measurements
- ED Care management consult
- Asthma education - compliance, identifying triggers, MDI teaching, smoking cessation

**Persistent or worsening symptoms < 48 hr L.O.S.**

- Increase frequency of nebulizer treatments
- IV Magnesium sulfate
- IV steroids
- Consider continuous nebs/IM epinephrine and transfer to CCT for further stabilization if severe deterioration

***DISPOSITION***

**Home:**

- Major resolution of sob/wheezing
- Peak flow >70% of predicted/baseline or significant improvement from baseline
- Ambulating comfortably
- Ensured follow up (PMD or Asthma/Chest clinic)
- Medication prescribed
- Consider escalation of outpt controller meds using stepwise approach if already compliant

**Admission:**

- Clinical deterioration to severe asthma exacerbation or imminent respiratory failure

Last updated 7/1/2019

Authored by E.Tang MD

Revised by T. Ahmad MD, R. Balakrishnan MD, A. Cai, MD, and S. Brewster MD

# ED OBSERVATION UNIT: ASTHMA GUIDELINES

## NYC H+H KINGS COUNTY HOSPITAL CENTER

Table 1: Suggested Peak Flow Rate When Height and Baseline measurements are not available <sup>6</sup>

| Asthma Severity | Peak Flow (L/min) |           |
|-----------------|-------------------|-----------|
|                 | Men               | Women     |
| Mild            | >400              | > 300     |
| Moderate        | 250 - 399         | 200 - 299 |
| Severe          | 150 - 249         | 120 - 200 |
| Very Severe     | <150              | < 120     |

### Sources

1. National Heart, Lung and Blood Institute: Guidelines for the Diagnosis and Management of Asthma, Expert Panel Report 3. Bethesda: National Institutes of Health Aug 2007.
2. Papiris, Spyros et al. "Clinical review: severe asthma." *Critical care (London, England)* vol. 6,1 (2001): 30-44.
3. Hodder R, Loughheed MD, Rowe BH, FitzGerald JM, Kaplan AG, McIvor RA. Management of acute asthma in adults in the emergency department: nonventilatory management. *CMAJ*. 2010;182(2):E55–E67. doi:10.1503/cmaj.080072
4. Landry A, Foran M, Koyfman A. Does Noninvasive Positive-Pressure Ventilation Improve Outcomes in Severe Asthma Exacerbations? *Ann Emerg Med* 2013;62(6):594-596
5. Lim WJ, Mohammed Akram R, Carson KV, Mysore S, Labiszewski NA, Wedzicha JA, Rowe BH, Smith BJ. Non-invasive positive pressure ventilation for treatment of respiratory failure due to severe acute exacerbations of asthma. *Cochrane Database Syst Rev*. 2012 Dec 12;12:CD004360.
6. Tsai CL, Clark S, Camargo CA, Jr. Risk stratification for hospitalization in acute asthma: the CHOP classification tree. *Am J Emerg Med*. 2010;28(7):803-808.

Last updated 7/1/2019

Authored by E.Tang MD

Revised by T. Ahmad MD, R. Balakrishnan MD, A. Cai, MD, and S. Brewster MD