



Appendix 1:

Age Based Non-Accidental Trauma Evaluation Guidelines

< 6 Months of Age

- Social Work Consult
- ACS contacted
- Skeletal Survey
- Ophthalmology Evaluation
- Head CT
- CMP, Phosphate, Lipase, Ca 2+ 25-hydroxyvitamin D
- Abdominal CT if AST/ALT > 80
- Troponin for infants > 3 months if AST/ALT > 80, signs of chest trauma, or ill appearance
- CBC, PT, and PTT if bruising or Intracranial Hemorrhage

6 – 12 Months of Age

- Social Work Consult
- ACS contacted
- Skeletal Survey
- Ophthalmology Evaluation
- CMP, Phosphate, Lipase, Ca 2+ 25-hydroxyvitamin D
- Abdominal CT if AST/ALT > 80
- Troponin if AST/ALT > 80, signs of chest trauma, or ill appearance
- CBC, PT, and PTT if bruising or Intracranial Hemorrhage
- Head CT for facial bruising, abnormal neurological examination or other clinical concern

12 – 36 Months of Age

- Social Work Consult
- ACS contacted
- CMP, Phosphate, Lipase, Ca 2+ 25-hydroxyvitamin D
- Skeletal Survey for < 24 months
 - Strongly consider skeletal Survey is >24 months and severely injured
- Abdominal CT if AST/ALT > 80
- Troponin if AST/ALT > 80, signs of chest trauma, or ill appearance
- CBC, PT, and PTT if bruising or Intracranial Hemorrhage
- Head CT for facial bruising, abnormal neurological examination or other clinical concern